



THE UNIVERSITY *of* EDINBURGH

## Edinburgh Research Explorer

# Inflammatory pain reduces C fibre activity-dependent slowing in a sex dependent manner, amplifying nociceptive input to the spinal cord

### Citation for published version:

Dickie, AC, McCormick, B, Lukito, V, Wilson, KL & Torsney, C 2017, 'Inflammatory pain reduces C fibre activity-dependent slowing in a sex dependent manner, amplifying nociceptive input to the spinal cord', *Journal of Neuroscience*, vol. 37, no. 27, pp. 6488-6502. <https://doi.org/10.1523/JNEUROSCI.3816-16.2017>

### Digital Object Identifier (DOI):

[10.1523/JNEUROSCI.3816-16.2017](https://doi.org/10.1523/JNEUROSCI.3816-16.2017)

### Link:

[Link to publication record in Edinburgh Research Explorer](#)

### Document Version:

Peer reviewed version

### Published In:

Journal of Neuroscience

### General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

### Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact [openaccess@ed.ac.uk](mailto:openaccess@ed.ac.uk) providing details, and we will remove access to the work immediately and investigate your claim.



Inflammatory pain reduces C fibre activity-dependent slowing in a sex dependent manner, amplifying nociceptive input to the spinal cord

Abbreviate title: Sex dependent C fibre ADS in inflammatory pain

Allen C Dickie<sup>1</sup>, Barry McCormick<sup>1</sup>, Veny Lukito<sup>1</sup>, Kirsten L Wilson<sup>1</sup> and Carole Torsney<sup>1</sup>

1. Centre for Integrative Physiology, Edinburgh Medical School: Biomedical Sciences, The University of Edinburgh, Edinburgh, EH8 9XD

Corresponding author:

Dr Carole Torsney, Centre for Integrative Physiology, Edinburgh Medical School: Biomedical Sciences, The University of Edinburgh, Hugh Robson Building, George Square, Edinburgh EH8 9XD. Email: [carole.torsney@ed.ac.uk](mailto:carole.torsney@ed.ac.uk)

# pages = 41 # figures = 8

# words: abstract (250) introduction (627) discussion (1500)

Conflict of interest: The authors declare no competing financial interests

Acknowledgements: This work was supported by a PhD scholarship and Centenary Award from the College of Medicine and Veterinary Medicine, The University of Edinburgh, both funded by the Medical Research Council. Allen C Dickie current affiliation: Spinal Cord Group, Institute of Neuroscience and Psychology, University of Glasgow. Barry McCormick current affiliation: MRC Centre for Inflammation Research, Queen's Medical Research Institute, University of Edinburgh. We thank Professor Mayank Dutia for helpful discussion.

## 28 Abstract

29 C fibres display activity-dependent slowing (ADS), whereby repetitive stimulation  
30 ( $\geq 1$ Hz) results in a progressive slowing of action potential conduction velocity, which  
31 manifests as a progressive increase in response latency. However, the impact of  
32 ADS upon spinal pain processing has not been explored nor whether ADS is altered  
33 in inflammatory pain conditions. To investigate, compound action potentials were  
34 made, from dorsal roots isolated from rats with or without complete Freund's  
35 adjuvant (CFA) hindpaw inflammation, in response to electrical stimulus trains. CFA  
36 inflammation significantly reduced C fibre ADS at 1 and 2Hz stimulation rates.  
37 Whole-cell patch-clamp recordings in the spinal cord slice preparation with attached  
38 dorsal roots, also demonstrated that CFA inflammation reduced ADS in the  
39 monosynaptic C fibre input to lamina I neurokinin 1 receptor expressing neurons (1-  
40 10Hz stimulus trains) without altering the incidence of synaptic response failures.  
41 When analysed by sex it was revealed that females display a more pronounced ADS  
42 that is reduced by CFA inflammation to a level comparable with males. Cumulative  
43 ventral root potentials evoked by long and short dorsal root stimulation lengths, to  
44 maximise and minimise the impact of ADS, respectively, demonstrated that reducing  
45 ADS facilitates spinal summation and this was also sex-dependent. This finding  
46 correlated with the behavioural observation of increased noxious thermal thresholds  
47 and enhanced inflammatory thermal hypersensitivity in females. We propose that  
48 sex/inflammation dependent regulation of C fibre ADS can, by controlling the  
49 temporal relay of nociceptive inputs influence the spinal summation of nociceptive  
50 signals contributing to sex/inflammation dependent differences in pain sensitivity.

## 51 Significance Statement

52 The intensity of a noxious stimulus is encoded by the frequency of action potentials  
53 relayed by nociceptive C fibres to the spinal cord. C fibres conduct successive action  
54 potentials at progressively slower speeds but the impact of this activity-dependent  
55 slowing (ADS) is unknown. Here we demonstrate that ADS is more prevalent in  
56 females than males and is reduced in an inflammatory pain model in females only.  
57 We also demonstrate a progressive delay of C fibre monosynaptic transmission to  
58 the spinal cord that is similarly sex and inflammation dependent. Experimentally  
59 manipulating ADS strongly influences spinal summation consistent with sex  
60 differences in behavioural pain thresholds. This suggests that ADS provides a  
61 peripheral mechanism that can regulate spinal nociceptive processing and pain  
62 sensation.

63

## 64 Introduction

65 The basic currency of communication in the nervous system is the action potential.  
66 As the action potential is an all-or-none event, information is coded by the number of  
67 action potentials and the time intervals between them. In the 1920's Edgar Adrian  
68 demonstrated that the intensity of a sensation is coded by the firing frequency in  
69 afferent nerve fibres in the somatosensory system (Adrian, 1926; Adrian and  
70 Zotterman, 1926a, b). Subsequently, microneurography studies in humans have  
71 directly demonstrated that the firing frequency in C fibre nociceptors encodes pain  
72 intensity (Torebjork et al., 1984; Yarnitsky and Ochoa, 1990). The more noxious the  
73 stimulus the shorter the intervals between successive action potentials in nociceptive  
74 C fibres and the more intense the pain experienced.

Nociceptive C fibres display *activity-dependent slowing* (ADS), whereby repetitive stimulation results in a progressive slowing of action potential conduction velocity, which manifests as a progressive increase in response latency in both human (Serra et al., 1999; Weidner et al., 1999) and animal studies (Thalhammer et al., 1994; Gee et al., 1996). Subtypes of C fibres display this phenomenon to differing degrees such that ADS can be used to functionally classify C fibre subtypes (Thalhammer et al., 1994; Serra et al., 1999; Weidner et al., 1999). ADS occurs in a frequency and length-dependent manner, with greater ADS observed at higher frequencies (Thalhammer et al., 1994; Gee et al., 1996; Serra et al., 1999; Weidner et al., 1999) and over longer lengths (Schmelz et al., 1995; Zhu et al., 2009). This progressive slowing of conduction velocity presumably regulates the intervals between successive action potentials reaching the spinal cord which could influence central pain processing and pain sensation, however, this has not been investigated.

ADS involves voltage-gated sodium (Nav) channels (De Col et al., 2008; Obreja et al., 2012) likely Nav1.7 and Nav1.8 (Baker and Waxman, 2012; Petersson et al., 2014; Tigerholm et al., 2014; Hoffmann et al., 2016) and is constrained by hyperpolarisation-activated cyclic nucleotide-gated (HCN) channels (Takigawa et al., 1998; Zhu et al., 2009; Mazo et al., 2013). These channels are known to be regulated in inflammatory pain (Emery et al., 2012; Weng et al., 2012; Rahman and Dickenson, 2013; Waxman and Zamponi, 2014) suggesting that C fibre ADS may be altered in inflammatory pain. Specifically, CFA inflammation increases both hyperpolarisation-activated current ( $I_h$ ) and HCN2 expression levels in C fibre nociceptors (Papp et al., 2010; Acosta et al., 2012; Weng et al., 2012) and alters  $I_h$  activation properties (Djouhri et al., 2015). Furthermore, deletion of HCN2 in Nav1.8 expressing neurons, which are mainly C fibre nociceptors limits inflammatory thermal

hyperalgesia (Emery et al., 2011) and peripheral block of HCN channels attenuates inflammatory pain (Young et al., 2014). There is also a loss of inflammatory pain phenotype in Nav1.7 and Nav1.8 knockout mice and in mice in which Nav1.7 was selectively deleted in Nav1.8 expressing neurons (Akopian et al., 1999; Nassar et al., 2004; Nassar et al., 2005). Furthermore, Nav1.7 and Nav1.8 channel expression is increased in CFA inflammation (Coggeshall et al., 2004; Gould et al., 2004; Liang et al., 2013) and selective blockers of either channel reduce inflammatory pain (McGowan et al., 2009; Zhang et al., 2010; Bregman et al., 2011; Yang et al., 2013; Lee et al., 2014; Payne et al., 2015).

The aim of this study was therefore to determine whether C fibre ADS is altered in the CFA inflammation model and the impact upon temporal relay of nociceptive input to the spinal cord. Given increasing awareness of sex differences in pain sensitivity and injury-induced hypersensitivity (Mogil and Bailey, 2010; Mogil, 2012; Bartley and Fillingim, 2013) this was also investigated in both sexes. Furthermore, the impact of ADS upon spinal summation and output was explored using electrophysiological and behavioural analysis of the nociceptive flexion withdrawal reflex.

## Materials and Methods

### *Animals*

All experiments were approved by the University of Edinburgh Ethical Review Committee and were performed in accordance with the UK Animals (Scientific Procedures) Act 1986. Sprague Dawley rats of both sexes (the University of Edinburgh Biological Research Resources, Edinburgh UK), were used in all

experiments. Animals were housed in cages at 21°C and 55% relative humidity, with a 12 h light-dark cycle, food and water were provided *ad libitum*.

#### *Inflammatory pain model*

To induce peripheral inflammation, juvenile rats received an intraplantar injection of complete Freund's adjuvant (CFA, 0.5 mg/ml saline) into the left hindpaw (1µl/g body weight) under isoflurane anaesthesia, at ~postnatal day (P)18, 2-5 days prior to patch-clamp or compound action potential electrophysiological recording at ~P21. This procedure results in persistent peripheral hindpaw inflammation and behavioural hypersensitivity in rats of this age (Torsney, 2011). Control rats were untreated.

#### *Isolated dorsal root preparation*

Isolated dorsal roots were prepared as described previously (Torsney, 2011; Dickie and Torsney, 2014). Briefly, naïve untreated (control) or CFA treated rats (~P21) were decapitated under isoflurane anaesthesia and spinal cords, with attached dorsal roots, were removed in ice-cold dissection solution. Lumbar (L4/L5) dorsal roots (left side only, CFA treated) were cut near the dorsal root entry zone and their dorsal root ganglia were removed, before being placed in 36–37°C oxygenated recovery solution for 1 hour. Roots were transferred to the recording chamber of an upright microscope (Zeiss) and were perfused with oxygenated Krebs' solution (1-2ml/min) at room temperature. The 95% O<sub>2</sub>/5% CO<sub>2</sub> saturated Krebs' solution contained (in mM): 125 NaCl, 2.5 KCl, 1.25 NaH<sub>2</sub>PO<sub>4</sub>, 26 NaHCO<sub>3</sub>, 25 glucose, 1 MgCl<sub>2</sub>, and 2 CaCl<sub>2</sub>, pH 7.4. Recovery solution was identical to Krebs' apart from 1.5mM CaCl<sub>2</sub> plus 6mM MgCl<sub>2</sub>. Dissection solution was the same as recovery, but with 1mM kynurenic acid.

147 *Compound action potential (CAP) recording*

148 Two glass suction electrodes were used, one for electrical orthodromic stimulation  
149 and the second for recording compound action potentials. Dorsal roots were  
150 stimulated 10 times at 0.2Hz (0.1ms duration), with an ISO-flex stimulus isolator  
151 (A.M.P.I. Intracel) at (in  $\mu\text{A}$ ): 1, 2, 3, 4, 5, 7.5, 10, 15, 20, 25, then in 10 $\mu\text{A}$  steps  
152 between 30 and 100 $\mu\text{A}$ , then in 50 $\mu\text{A}$  steps between 150 and 500 $\mu\text{A}$  (Torsney, 2011;  
153 Dickie and Torsney, 2014). An 0.1ms pulse width was chosen to replicate the  
154 electrical stimuli previously established to activate the different afferent fibre types in  
155 this age of rat (Nakatsuka et al., 2000). However, the possibility of an  
156 underestimation of the C fibre contribution cannot be excluded given that longer  
157 pulse widths have also been employed to stimulate C fibre inputs (Baba et al., 1999).  
158 The main components of the compound action potentials were differentiated as A $\beta$ ,  
159 A $\delta$  and C fibre, on the basis of activation threshold and conduction velocity, each  
160 displaying a characteristic triphasic (positive-negative-positive) response. Data were  
161 acquired and recorded using a Cygnus ER-1 differential amplifier (Cygnus  
162 Technologies Inc.) and pClamp 10 software (Molecular Devices). Data were filtered  
163 at 10kHz and sampled at 50kHz.

164 Activation threshold was defined as the lowest stimulation intensity at which the  
165 negative component of the triphasic response was clearly identifiable. The amplitude  
166 of each component was calculated by measuring the distance between the negative  
167 and second positive peak. The conduction velocity was calculated based on the  
168 latency to the negative peak at 20, 100 and 500 $\mu\text{A}$  for A $\beta$ , A $\delta$  and C components,  
169 respectively.

170 To assess ADS, dorsal roots were stimulated 16 or 40 times (500 $\mu\text{A}$  intensity, 0.1ms  
171 duration) at frequencies of 1 or 2Hz. For each stimulus, the latency between the



stimulus artefact and the negative peak of the triphasic response was measured and the change in latency from stimulus 1 calculated. In some cases the width of the C fibre component (positive peak to positive peak) was additionally measured and the change in width from stimulus 1 calculated. To negate any influence of varying dorsal root length, the latency/width change was normalised to the length of root stimulated, measured as the distance between the stimulating and recording electrodes. In a subset of recordings the stimulating electrode was first placed close to the distal end of the dorsal root (long stimulation length) and then placed closer to the recording electrode (short stimulation length). By subtracting the latency change values for 'short stimulation length' from 'long stimulation length' the latency change solely attributable to conduction velocity slowing, independent of action potential initiation was calculated.

#### *Spinal cord slice preparation*

Spinal cords with attached dorsal roots, from which dorsal root ganglia were removed, were obtained from control or CFA treated (~P21) rats, as described above. The lumbar (L4/L5) segment was embedded in an agarose block and 350µm slices, with attached dorsal roots (left side only, CFA treated), were cut. Slices were placed in oxygenated recovery solution at 36–37°C for 1 hour and were then incubated at room temperature for 30 mins with 35nM tetramethylrhodamine conjugated substance P (TMR-SP), as described previously (Labrakakis and MacDermott, 2003; Torsney, 2011; Dickie and Torsney, 2014). Slices were allowed to recover for a further 1 hour at room temperature before being transferred to the recording chamber of an upright microscope (Zeiss), equipped with fluorescence for the identification of TMR-SP labelled (TMR-SP+) neurons and infrared-differential

196 interference contrast (IR-DIC) for electrophysiological recordings, and were  
197 continually perfused with oxygenated Krebs (1-2ml/min) at room temperature.

#### 198 *Patch-clamp recording*

199 Whole-cell patch-clamp recordings (holding potential -70mV) were made from TMR-  
200 SP+ neurons in the lamina I region of the dorsal horn. The intracellular solution used  
201 was composed of (in mM); 120 Cs-methylsulfonate, 10 Na-methylsulfonate, 10  
202 EGTA, 1 CaCl<sub>2</sub>, 10 HEPES, 5 QX-314-Cl [N-(2,6-  
203 Dimethylphenylcarbamoylmethyl)triethylammonium chloride] and 2 Mg<sup>2+</sup>-ATP, pH  
204 adjusted to 7.2 with CsOH, osmolarity 290mOsm and junction potential was  
205 corrected prior to recording. Additionally, 1μM Alexa Fluor 488 hydrazide was  
206 included in the recording pipette. Data were recorded and acquired with an Axopatch  
207 200B amplifier and pClamp 10 software (Molecular Devices). Data were filtered at  
208 5kHz and digitised at 10kHz.

209

210 Monosynaptic primary afferent input to lamina I NK1R+ neurons was identified as  
211 described previously (Torsney and MacDermott, 2006; Torsney, 2011; Dickie and  
212 Torsney, 2014). eEPSCs were recorded in response to low frequency (0.05Hz)  
213 dorsal root stimulation (3 times) at intensities of 20, 100 and 500μA (0.1ms stimulus  
214 duration) to activate Aβ, Aδ and C fibre inputs, respectively, using an ISO-Flex  
215 stimulus isolator. To characterise an input as monosynaptic or polysynaptic, dorsal  
216 roots were stimulated (20 times) at the following intensities and frequencies; Aβ  
217 20μA / 20Hz; Aδ 100μA / 2Hz; C 500μA / 1Hz. A-fibre responses were considered  
218 monosynaptic if they displayed no synaptic failures and a stable latency (≤2ms),

while C fibre inputs were considered monosynaptic if they displayed no synaptic failures, regardless of whether there was latency variability (Nakatsuka et al., 2000).

To assess whether monosynaptic A $\delta$  or monosynaptic C fibre input to lamina I NK1R+ neurons displayed ADS in response to repetitive stimulation, eEPSCs were recorded in response to trains of 16 stimuli delivered at 1 or 2Hz (A $\delta$  and C) or trains of 40 stimuli at 2, 5 or 10Hz (C only), at intensities of 100 (A $\delta$ ) or 500 $\mu$ A (C). The latency of each eEPSC was measured as the time between the stimulus artefact and the onset of the monosynaptic response and the change in latency from stimulus 1 calculated. These latency change data were also normalised to dorsal root length, measured as the distance between the stimulating electrode and the dorsal root entry zone, to account for variations in the length of dorsal root stimulated.

#### *Dorsal root – ventral root potential (DR-VRP) recording*

Control rats (~P10) were decapitated under isoflurane anaesthesia and spinal cords, with attached dorsal and ventral roots, were removed in ice-cold dissection solution. Hemisected lumbar spinal cord with only L4/L5 dorsal and ventral roots left attached were prepared (Otsuka and Konishi, 1974) and transferred to a recording chamber perfused with oxygenated Krebs' solution (>2ml/min) at room temperature. Two glass suction electrodes were used, one for electrical stimulation of the dorsal root and the second for recording ventral root potentials. The stimulating electrode was first placed close to the distal end of the dorsal root (long stimulation length/more ADS) and then placed closer to the spinal cord (short stimulation length/less ADS) to assess the impact of length-dependent ADS upon spinal summation. The dorsal root was stimulated 40 times at 2, 5 and 10Hz (500 $\mu$ A intensity, 0.1ms duration) at both

sites and the cumulative ventral root potential recorded using a close fitting glass electrode placed on the ventral root close to the ventral horn (Thompson et al., 1994). Data were acquired and recorded using a Cygnus ER-1 differential amplifier (Cygnus Technologies Inc.) and pClamp 10 software (Molecular Devices). Data were filtered at 10kHz and sampled at 50kHz.

#### *Sensory testing*

Prior to and 2-5 days following intraplantar CFA injection (as detailed above) hindpaw swelling, mechanical and thermal sensitivity were measured. Hindpaw swelling was assessed by measuring the thickness of the dorsal-ventral paw using Vernier calipers. Following habituation on an elevated mesh platform the mechanical threshold of the nociceptive flexion withdrawal reflex was determined with von Frey filaments (Stoelting) applied to the mid-plantar surface of the hindpaw using the up-down method (Chaplan et al., 1994). After habituation to the Hargreaves apparatus, radiant heat was applied to the mid-plantar surface of the hindpaw (x3 stimuli per hindpaw to calculate average) to determine the noxious thermal withdrawal latency.

#### *Statistical analysis*

Area under the curve (AUC) analysis was used to compare ADS in both CAP and patch-clamp recordings. Group comparisons, for both electrophysiology and behavioural data, were carried out using 2-way ANOVA with or without repeated measures analysis as appropriate and was followed by Sidak's multiple comparisons test or Tukey's multiple comparisons test, respectively, if an interaction between factors was observed. Averaged data are represented as mean  $\pm$  SE.

## Materials

All chemicals were obtained from Sigma, except; TMR-SP (Enzo Life Sciences), Alexa Fluor 488 hydrazide (Molecular Probes), QX-314-Cl (Alomone Labs) and NBQX (Tocris Bioscience).

## Results

### *CFA inflammation reduces C fibre ADS in isolated dorsal roots*

To assess the impact of CFA inflammation on C fibre ADS, dorsal roots isolated from control or CFA treated rats were repetitively stimulated and the response latencies of A $\beta$ , A $\delta$  and C fibre compound action potentials were measured. The A $\beta$ , A $\delta$  and C fibre afferent components were identified on the basis of conduction velocity and activation threshold (Fig. 1A). CFA inflammation did not alter threshold stimulus intensity ( $P=0.394$ : A $\beta$  control ( $n=10$ ),  $7.6\pm0.80$   $\mu$ A; A $\beta$  CFA ( $n=12$ ),  $5.9\pm0.60$   $\mu$ A; A $\delta$  control ( $n=11$ ),  $38.6\pm3.0$   $\mu$ A; A $\delta$  CFA ( $n=13$ ),  $34.6\pm2.8$   $\mu$ A; C control ( $n=11$ ),  $240.9\pm14.8$   $\mu$ A; C CFA ( $n=13$ )  $226.9\pm15.6$   $\mu$ A) conduction velocity ( $P=0.418$ : A $\beta$  control,  $4.6\pm0.50$  m/s; A $\beta$  CFA,  $5.0\pm0.30$  m/s; A $\delta$  control,  $0.9\pm0.1$  m/s; A $\delta$  CFA,  $0.7\pm0.1$  m/s; C control,  $0.2\pm0.01$  m/s; C CFA  $0.2\pm0.01$  m/s) or amplitude ( $P=0.091$ : A $\beta$  control,  $1.7\pm0.2$  mV; A $\beta$  CFA,  $2.4\pm0.40$  mV; A $\delta$  control,  $0.1\pm0.02$  mV; A $\delta$  CFA,  $0.1\pm0.02$  mV; C control,  $0.1\pm0.02$  mV; C CFA  $0.1\pm0.03$  mV), as previous studies have demonstrated in both adult (Baba et al., 1999; Nakatsuka et al., 2000) and similarly aged juvenile rats (Torsney, 2011).

Repetitive stimulation of isolated dorsal roots produced a negligible reduction (speeding) in the latency of the A $\beta$  fibre response (Fig. 1Bi,Ci) and a marginal

increase (slowing) in the A $\delta$  fibre response latency (Fig. 1Bii,Cii). CFA inflammation did not alter these observations in A fibres (A $\beta$ :  $P=0.693$ ; A $\delta$ :  $P=0.451$ ). In contrast, repetitive stimulation resulted in a clear progressive increase in C fibre response latency (Fig. 1Biii), that was confirmed to be frequency-dependent, with 2Hz stimulation resulting in greater ADS than 1Hz stimulation ( $P<0.0001$ , Fig. 1Ciii). Notably, this C fibre ADS was significantly reduced by CFA inflammation, independent of stimulation frequency ( $P<0.0001$ , Fig. 1Biii, Ciii).

To address the possibility that the observed CFA effect may reflect altered action potential initiation time rather than altered conduction velocity slowing, compound action potentials were recorded with the stimulating electrode placed at two different positions on an individual dorsal root (Fig. 1Di). Subtraction of the C fibre latency values recorded following stimulation at position 2 (short stimulation length) from those stimulated at position 1 (long stimulation length) eliminates the contribution of action potential initiation, leaving only conduction time (between the 2 stimulation sites). This also revealed a progressive increase in C fibre response latency (Fig. Dii) that was reduced by CFA inflammation independent of stimulus frequency ( $P=0.009$  Fig. 1Dii,iii). This suggests that altered action potential initiation time is not therefore likely to account for the observed CFA inflammation dependent reduction in C fibre ADS. Furthermore, the demonstration that subtracting the short from long stimulation length results in a progressive increase in the C fibre response latency confirms the length dependency of the phenomenon.

#### *CFA inflammation reduces ADS in monosynaptic C fibre input to lamina I NK1R+ neurons*

To explore the spinal impact of CFA reduced C fibre ADS, whole-cell patch-clamp recordings were made from lamina I NK1R+ neurons, that are likely projection

neurons (Marshall et al., 1996; Todd et al., 2000; Spike et al., 2003; Al-Khater et al., 2008) that receive monosynaptic input from both C and A $\delta$  fibres (Torsney and MacDermott, 2006; Torsney, 2011; Peirs et al., 2015). Comparison of the synaptic transmission of these inputs, with their distinct afferent temporal relays (Fig. 1) will provide insight into the central impact of ADS and its regulation by CFA. Lamina I NK1R+ neurons were pre-identified using TMR-SP, which is not expected to alter recorded synaptic activity (Tong and MacDermott, 2006), and has been employed previously (Tong and MacDermott, 2006; Torsney and MacDermott, 2006; Torsney, 2011; Dickie and Torsney, 2014; Peirs et al., 2015).

Monosynaptic A $\delta$  fibre eEPSCs were recorded, in spinal cord slices from control and CFA treated rats (Fig. 2A), in response to dorsal root stimulation at frequencies of 1 and 2Hz. This resulted in a small, frequency dependent ( $P=0.016$ ), progressive increase in the eEPSC latency, that was unaffected by CFA inflammation ( $P=0.570$ , Fig. 2B,C).

In contrast, repetitive stimulation of monosynaptic C fibre input to lamina I NK1R+ neurons resulted in a progressive increase in response latency, that was not altered by stimulation frequency ( $P=0.521$ ) but was markedly reduced by CFA inflammation ( $P=0.013$ , Fig. 3) similar to the population C fibre CAP recordings. Notably, the initial conduction velocity of monosynaptic C fibre input to lamina I NK1R+ neurons was not altered by CFA inflammation ( $p=0.764$ , Mann-Whitney, data not shown), as demonstrated previously (Torsney, 2011). CFA inflammation also did not alter the initial peak amplitude of C fibre eEPSCs ( $p=0.568$ , Mann Whitney, data not shown), as reported previously (Torsney, 2011; Dickie and Torsney, 2014) or the eEPSC amplitude observed during repetitive stimulation ( $P=0.178$ , 2-way ANOVA, data not shown). In summary, CFA inflammation does not alter the baseline CV of

monosynaptic C fibre inputs or their eEPSC peak amplitudes in lamina I NK1R+ neurons but it significantly reduces the progressive delay in synaptic transmission observed between C fibres and lamina I NK1R+ neurons.

*CFA inflammation reduces both the average and range of ADS within the C fibre population*

Longer stimulus trains and higher frequencies produce a more pronounced C fibre ADS that is associated with the occurrence of conduction failures (Thalhammer et al., 1994; Nakatsuka et al., 2000; Zhu et al., 2009) the prevalence of which can be altered in pain models (Sun et al., 2012; Wang et al., 2016a) which could additionally impact upon spinal pain processing. Therefore, to investigate the potential impact of conduction failures, longer stimulus trains and higher frequencies were employed.

In CAP recordings stimulating isolated dorsal roots with trains of 40 stimuli delivered at 1 and 2Hz resulted in a progressive increase in the C fibre response latency, which was frequency-dependent ( $p < 0.0001$ ) and was significantly reduced by CFA inflammation ( $p = 0.005$ , Fig. 4A). It was not possible to employ higher frequency stimulation in the CAP recordings as the C fibre component of the compound action potential diminishes substantially during repetitive stimulation, presumably due to conduction failures, and as such could only be reliably quantified at 1 and 2 Hz. The change in C fibre response latency reflects the change in average conduction velocity of the population C fibre response. Given that different C fibres display different degrees of ADS (Thalhammer et al., 1994; Serra et al., 1999; Weidner et al., 1999) the change in width of the C fibre response was also measured (Fig. 4B), as this will reflect the change in range of conduction velocities present within the population and may therefore be a more informative measure of ADS across the entire C fibre population. 1 and 2Hz stimulation resulted in a progressive increase in the C fibre response width which was also regulated in a frequency and CFA



inflammation dependent manner (both  $p < 0.0001$ , Fig. 4C). Notably, the initial width of the C fibre response was not significantly different between control and CFA tissue ( $p = 0.27$ , unpaired t-test, data not shown).

*CFA inflammation limits the progressive delay in synaptic transmission between C fibres and lamina I NK1R+ neurons without altering synaptic response failures.*

In eEPSC recordings repetitive stimulation (x40) of the monosynaptic C fibre input to lamina I NK1R+ neurons at 2, 5 and 10Hz, resulted in a frequency-dependent ADS ( $P = 0.039$ ) that was reduced by CFA inflammation ( $p < 0.0001$ , Fig. 5A). Plotting the percentage of lamina I NK1R+ neurons displaying synaptic response failures per stimulus number reveals a progressive increase in the number of failures (Fig. 5Bi-iii). This progressive increase in synaptic response failures is, as expected, frequency dependent, with the total no of failures per lamina I NK1R+ neuron increasing with stimulation frequency ( $p < 0.0001$ ). However, the total no of synaptic response failures was not affected by CFA inflammation ( $p = 0.971$ , Fig. 5Biv).

*C fibre ADS is regulated in a sex and inflammation dependent manner*

To determine whether there is a sex difference in the ADS phenomenon CAP (Fig. 6A-C) and eEPSC (Fig. 6D-F) datasets were analysed for females and males separately. In CAP recordings from isolated dorsal roots a progressive increase in the latency and width of the population C fibre response was observed in both females (Fig. 6Ai,ii) and males (Fig. 6Bi,ii). AUC analysis reveals a significant interaction between sex and CFA inflammation for both latency change (Fig. 6Ci  $p = 0.037$ ) and width change (Fig. 6Cii  $p = 0.003$ ). ADS is more pronounced in control females and CFA inflammation reduces ADS, in females only, to a level that is now comparable with males for both latency change (CFA: female  $p = 0.031$ ; male

p=0.998) and width change measures (CFA: female  $p < 0.0001$ ; male  $p = 0.525$ ). Initial CV and CAP width were not altered in a sex or CFA inflammation dependent manner (CV: sex  $p = 0.853$ , CFA  $p = 0.120$ ; width: sex  $p = 0.482$ , CFA  $p = 0.307$ , 2-way ANOVA data not shown).

Similarly, patch-clamp recordings from lamina I NK1R+ neurons show a progressive increase in the latency of monosynaptic C fibre evoked eEPSCs in both females (Fig. 6D) and males (Fig. 6E). Likewise AUC analysis reveals a significant interaction between sex and CFA inflammation (Fig. 6F  $p = 0.034$ ). ADS is more pronounced in control females and CFA inflammation reduces ADS, in females only, to a level that is now comparable with males (CFA: female  $p = 0.015$ ; male  $p = 0.996$ ). However, baseline CV of monosynaptic C fibre inputs was not altered in a sex or CFA inflammation dependent manner (sex  $p = 0.873$ ; CFA  $p = 0.923$ , data not shown).

#### *Limiting ADS facilitates spinal summation*

Our observations demonstrate that ADS results in a progressive delay in the synaptic transmission of C fibre input to individual spinal neurons. Moreover, given that repetitive C fibre stimulation results in a progressive increase in C fibre CAP width, that reflects the range of CVs within the population, ADS also likely reduces the temporal coincidence of population C fibre input at a spinal network level.

Together, these findings suggest that ADS limits temporal summation of C fibre evoked synaptic activity at a spinal level. Thereby, reducing C fibre ADS should enhance spinal summation. To test this hypothesis we took advantage of the length dependency of ADS (Schmelz et al., 1995; Zhu et al., 2009) and assessed the impact of long (increased ADS) versus short (decreased ADS) root stimulation lengths upon spinal summation by recording cumulative dorsal root evoked ventral root potentials in a hemisectioned spinal cord preparation (Fig. 7A). Using this

preparation we are essentially able to assess the impact of ADS upon the circuitry underlying the nociceptive flexion withdrawal reflex.

Repetitive stimulation of dorsal roots evoked a cumulative ventral root potential that is blocked, as expected (Thompson et al., 1992) in the presence of the AMPA receptor antagonist NBQX (10 $\mu$ M) and NMDA receptor antagonist APV (50 $\mu$ M, Fig. 7B). This spinal summation is frequency dependent (Fig. 7Ci), as previously reported (Thompson et al., 1992). Interestingly, reducing the stimulation length strongly facilitated the cumulative ventral root potential at 2Hz stimulation rates (Fig. 7Cii). To quantify the degree of summation the amplitude of the cumulative ventral root potential was measured 500ms following the last stimulus artefact and normalised to the amplitude measured at 500ms following a single stimulus at long or short root stimulation length to account for any potential difference in response amplitude between stimulation lengths. Repeated measures ANOVA revealed a significant interaction ( $p=0.022$ ) between frequency ( $p=0.039$ ) and length ( $p=0.003$ ) dependent effects and post tests revealed a significant difference between long and short lengths at 2Hz stimulation rates only (Fig. 7Di, 2Hz  $p<0.01$ ). Given the more pronounced C-fibre ADS in females we predicted a greater change in facilitation of spinal summation in females versus males. Indeed, there is a greater difference in the % cumulative depolarisation between long and short stimulation length in females at the 2Hz stimulation rate (Fig 7Dii,  $p=0.028$ ).

To estimate the degree of change in ADS resulting from the average ~50% reduction in root stimulation length in the DR-VRP recordings and thus the likely relevance of the findings with respect to our sex/inflammation dependent changes in ADS we revisited the x2 length stimulation CAP recordings displayed in Figure 1D. In the CAP recordings a comparable reduction in stimulation length (~60%) reduced ADS

by ~20% as measured using AUC analysis of latency change (unpaired t-test  $p=0.007$ , data not shown). Therefore in these DR-VRP recordings the short stimulation length likely underestimates the average ~30% reduction in ADS compared with the control female grouping (also assessed using AUC analysis of latency change, Fig. 6Ci) underscoring the importance of these findings.

*Females display elevated noxious thermal thresholds and enhanced inflammatory thermal hyperalgesia*

The demonstration that females display more pronounced ADS that is reduced by CFA inflammation to a level that is comparable with males (control and CFA) along with the observation that limiting ADS facilitates cumulative dorsal root evoked ventral root potentials, the following predictions can be made regarding the nociceptive flexion withdrawal reflex: 1) control females should have higher pain thresholds than control males; 2) if other underlying inflammatory pain mechanisms are not sex-dependent, then CFA inflammation reduced pain thresholds should be similar between males and females and 3) because these predictions imply a greater reduction in pain threshold in females vs males, females should display enhanced inflammatory hypersensitivity.

Figure 8A-C displays behavioural data obtained in response to noxious radiant heat stimuli in the CFA inflammation model that supports these predictions. Females show increased inflammatory thermal hypersensitivity (Fig. 8B,  $p=0.032$ ) that, importantly, does not appear to be due to an increased inflammatory response because CFA induced paw swelling is not significantly different between sexes (Fig. 8D,  $p=0.513$ ). It instead reflects a larger reduction in thermal pain thresholds from an elevated baseline level as can be observed by comparing contralateral male and female withdrawal latencies (Fig. 8A). Notably, AUC analysis of thermal withdrawal

latency values (2-5d post CFA) reveals a significant interaction ( $p=0.044$ ) between sex and hindpaw (ipsilateral/contralateral) with post-tests revealing significantly higher control/contralateral hindpaw values in females ( $p=0.022$ ). The mechanical threshold of the flexion withdrawal reflex was assessed using von Frey monofilaments (Fig. 8E). AUC analysis reveals no significant interaction between sex and hindpaw (repeated measures ANOVA  $p=0.180$ , data not shown). Furthermore, inflammatory mechanical hypersensitivity was not altered in a sex-dependent manner (Fig. 8F,  $p=0.548$ ).

## Discussion

We found that C fibre ADS is more pronounced in females than males and is reduced by CFA inflammation in females only. This alters the timing of synaptic transmission of monosynaptic C fibre input to lamina I NK1R+ neurons. Experimental manipulation of ADS demonstrates it can influence spinal summation consistent with observed sex differences in noxious thermal thresholds. We propose that ADS regulates nociceptive drive to central pain circuits.

### *Underlying mechanisms of ADS*

The underlying mechanism of ADS is not fully understood. It was initially proposed to result from  $\text{Na}^+\text{-K}^+\text{-ATPase}$  dependent membrane hyperpolarisation (Rang and Ritchie, 1968; Bostock and Grafe, 1985) but  $\text{Na}^+\text{-K}^+\text{-ATPase}$  blockade increased rather than reduced ADS (De Col et al., 2008). Pharmacological studies then suggested ADS reflects increased numbers of Nav channels entering a slow inactivated state (De Col et al., 2008; Obreja et al., 2012). However modelling suggested that while Nav channel slow inactivation contributes, ADS is most readily associated with an increase in intra-axonal  $\text{Na}^+$  concentration (Tigerholm et al.,

2014). Investigation of the differences between mechano-insensitive C fibres that display the greatest ADS versus mechano-sensitive C fibres that display minimal ADS (Weidner et al., 1999; Obreja et al., 2010) revealed that pronounced ADS was associated with more Nav1.8, less Nav1.7, more delayed rectifier potassium channel and less Na<sup>+</sup>/K<sup>+</sup> ATPase, a profile consistent with enhanced accumulation of intracellular Na<sup>+</sup> (Petersson et al., 2014). Notably, given the observed sex/inflammation dependent ADS these molecules are regulated by peripheral inflammation (Coggeshall et al., 2004; Gould et al., 2004; Zhang and Nicol, 2004; Liang et al., 2013; Rahman and Dickenson, 2013; Waxman and Zamponi, 2014; Wang et al., 2015) but, despite being key pharmaceutical targets, sex differences in their regulation in C fibre nociceptors have not been explored.

In addition to the aforementioned molecules implicated in ADS there are many other ion channels/receptors whose expression, location and/or function can be altered by inflammation (Gold and Gebhart, 2010) and may contribute to altered ADS. Furthermore, previous studies have reported CFA inflammation induced changes in C-fibre nociceptor excitability including increased CV combined with reduced electrical thresholds in guinea-pig (Djouhri and Lawson, 2001) not replicated in this or other rat studies (Baba et al., 1999; Nakatsuka et al., 2000; Torsney, 2011), altered action potential shape (Djouhri and Lawson, 1999; Zhang et al., 2012) and spontaneous firing (Djouhri et al., 2006; Xiao and Bennett, 2007; Matson et al., 2015). Moreover, estrogen exacerbates inflammation increased excitability of temporomandibular joint afferents (Flake et al., 2005) and there are sex differences in inflammatory sensitisation of dural afferents (Scheff and Gold, 2011).

511 *Evidence for altered ADS*

512 Consistent with our observed inflammation altered ADS, the inflammatory mediator  
513 NGF reduces ADS in the mechano-insensitive C fibres that normally display  
514 pronounced ADS (Obreja et al., 2011) and is associated with less Nav1.8, more  
515 Nav1.7, less delayed rectifier potassium channel and more Na<sup>+</sup>/K<sup>+</sup> ATPase  
516 (Petersson et al., 2014). Furthermore, there is reduced ADS in a diabetic  
517 neuropathic pain model (Wang et al., 2016b). In contrast, there is enhanced ADS in  
518 the spinal nerve ligation model of neuropathic pain (Shim et al., 2007) but these  
519 findings need to be considered within a partially denervated pain circuitry.  
520 Microneurography studies have also demonstrated altered ADS in chronic pain  
521 patients (Orstavik et al., 2003; Orstavik et al., 2006; Kleggetveit et al., 2012; Serra et  
522 al., 2014), interestingly, including patients with mutations in Nav1.7 (Namer et al.,  
523 2015) and Nav1.8 (Kist et al., 2016).

524 *Physiological relevance of ADS*

525 C fibre ADS is physiologically relevant because it occurs, not only following electrical  
526 stimulation, but also in response to natural stimulation of the skin (Thalhammer et al.,  
527 1994). Electrical stimulation induced ADS is useful as it can provide a readout of  
528 nociceptor excitability given the key roles of molecules implicated in ADS, such as  
529 Nav and HCN channels, in action potential generation and regulation of initial firing  
530 frequency in nociceptor terminals. It is not surprising therefore that altered ADS  
531 profiles are associated with changes in C fibre thresholds (Obreja et al., 2011), that  
532 ADS correlates with spontaneous firing (Kleggetveit et al., 2012) and is accompanied  
533 by a parallel increase in C fibre mechanical threshold (De Col et al., 2012).

534

Moreover, ADS is proposed to provide a memory trace of previous activity levels that can influence responses to subsequent inputs. Specifically, low level firing, comparable to spontaneous firing rates in inflammatory pain induces ADS that dynamically influences the response to higher frequency inputs such that they display reduced ADS or even speeding (Weidner et al., 2002). Importantly, in our data inflammation induced spontaneous C fibre firing will likely be absent, due to the lack of peripheral inflamed tissue, in our *ex vivo* preparations. Therefore our observations may well underestimate the degree of inflammation reduced ADS as our recorded data likely only reflects the inflammation altered expression levels of ion channels involved in ADS and not the dynamic regulation of ADS by ongoing C fibre activity.

Here we demonstrate, that in addition to ADS providing a nociceptor excitability readout and a dynamic memory, that the progressive slowing per se significantly influences the relay of nociceptive signals such that the firing frequency initiated in the periphery is not faithfully transmitted to the spinal cord. In individual spinal neurons there is a progressive delay in synaptic transmission of C fibre inputs and we also observe a progressive reduction in temporal coincidence of population C fibre input that we predicted should alter summation at a spinal network level. Decreasing stimulation length in DR-VRP recordings demonstrated that reducing ADS did indeed facilitate spinal summation. Interestingly, facilitation was only significant at the 2Hz stimulation rate and strikingly, increased summation to a level comparable with that observed at 10Hz. The lack of effect at higher frequencies probably reflects the increased incidence of synaptic failures with increasing stimulation frequency (see Figure 5B) likely resulting from frequency dependent C



fibre conduction failure (Nakatsuka et al., 2000; Zhu et al., 2009; Sun et al., 2012; Wang et al., 2016a) that will limit the maximal extent of spinal summation. Notably, spontaneous firing rates resulting from tissue inflammation are typically <1Hz (Djouhri et al., 2006; Xiao and Bennett, 2007; Matson et al., 2015) whereas noxious stimuli evoke firing in the 1-10Hz range (Lynn and Carpenter, 1982; Leem et al., 1993). We therefore propose, that reducing ADS thereby promoting temporal coincidence of nociceptive input, as a novel mechanism of hyperalgesia whereby lower intensity noxious inputs can be amplified but high intensity inputs are not further amplified due to transmission failures which may reflect an intrinsic self-inhibition mechanism to limit overdrive of the nociceptive pathways. Notably ADS was altered in the monosynaptic C fibre input to lamina I NK1R+ likely projection neurons that display tightly controlled spike timing-dependent plasticity (Li and Baccei, 2016). Therefore these changes in ADS may influence the involvement of these output neurons in synaptic plasticity (Ikeda et al., 2003; Ikeda et al., 2006) spinal supra-spinal loop activity (Suzuki et al., 2002) and chronic pain (Mantyh et al., 1997; Nichols et al., 1999). Finally promotion of closely timed nociceptive inputs appears to be a feature following injury with the recent demonstration of novel DRG neuronal co-activation following CFA inflammation that contributes to mechanical hyperalgesia (Kim et al., 2016).

The pronounced ADS in control females that was reduced by CFA to male levels, along with the observation that limiting ADS facilitates spinal summation, was consistent with behavioural observations of elevated noxious thermal latencies and enhanced inflammatory thermal hyperalgesia in females. While the enhanced hypersensitivity is consistent with sex differences in human studies the elevated

noxious thermal latencies are not (Mogil, 2012; Bartley and Fillingim, 2013). However, we studied spinal reflex behaviours in which the afferent input is a major component of the underlying neural circuitry whereas human subjective pain scoring additionally involves higher level cognitive processing. If differing degrees of ADS contributes to sex differences in pain thresholds, it is not surprising that this is specific for thermal vs mechanical sensitivity because the inflammatory mediator NGF can reduce ADS in mechano-insensitive C fibres that can respond to heat (Schmidt et al., 1995; Schmelz et al., 1997) without unmasking of mechanical sensitivity (Obreja et al., 2011). However, the lack of sex differences in mechanical sensitivity may reflect the use of von Frey hairs to identify mechanical thresholds rather than a noxious mechanical stimulus. Alternatively it may be that different peripheral mechanisms, DRG neuronal coupling and reduced ADS are employed to augment noxious mechanical and thermal inputs, respectively. It will be important to determine the extent to which ADS regulates the processing of noxious stimuli *in vivo* and also identify the mechanisms underlying sex dependent ADS and whether these are controlled by sex hormones or genetic factors.

It has been long established that the intensity of a sensation is encoded by afferent firing frequency. Here we propose that the firing frequency initiated in the periphery, is diminished en route to the CNS, by ADS in nociceptive C fibres and that inflammation/sex dependent regulation of ADS can thereby peripherally modulate spinal processing and pain sensation.

## References

609 Acosta C, McMullan S, Djouhri L, Gao L, Watkins R, Berry C, Dempsey K, Lawson SN (2012) HCN1 and  
610 HCN2 in Rat DRG neurons: levels in nociceptors and non-nociceptors, NT3-dependence and  
611 influence of CFA-induced skin inflammation on HCN2 and NT3 expression. PLoSOne  
612 7:e50442.

613 Adrian ED (1926) The impulses produced by sensory nerve-endings: Part 4. Impulses from Pain  
614 Receptors. J Physiol 62:33-51.

615 Adrian ED, Zotterman Y (1926a) The impulses produced by sensory nerve-endings: Part II. The  
616 response of a Single End-Organ. J Physiol 61:151-171.

617 Adrian ED, Zotterman Y (1926b) The impulses produced by sensory nerve endings: Part 3. Impulses  
618 set up by Touch and Pressure. J Physiol 61:465-483.

619 Akopian AN, Souslova V, England S, Okuse K, Ogata N, Ure J, Smith A, Kerr BJ, McMahon SB, Boyce S,  
620 Hill R, Stanfa LC, Dickenson AH, Wood JN (1999) The tetrodotoxin-resistant sodium channel  
621 SNS has a specialized function in pain pathways. NatNeurosci 2:541-548.

622 Al-Khater KM, Kerr R, Todd AJ (2008) A quantitative study of spinothalamic neurons in laminae I, III,  
623 and IV in lumbar and cervical segments of the rat spinal cord. JComp Neurol 511:1-18.

624 Baba H, Doubell TP, Woolf CJ (1999) Peripheral inflammation facilitates Abeta fiber-mediated  
625 synaptic input to the substantia gelatinosa of the adult rat spinal cord. JNeurosci 19:859-867.

626 Baker MD, Waxman SG (2012) Sodium channel slow inactivation and adaptation in C-fibres. J Physiol  
627 590:1513-1514.

628 Bartley EJ, Fillingim RB (2013) Sex differences in pain: a brief review of clinical and experimental  
629 findings. BrJAnaesth 111:52-58.

630 Bostock H, Grafe P (1985) Activity-dependent excitability changes in normal and demyelinated rat  
631 spinal root axons. J Physiol 365:239-257.

632 Bregman H et al. (2011) Identification of a potent, state-dependent inhibitor of Nav1.7 with oral  
633 efficacy in the formalin model of persistent pain. JMedChem 54:4427-4445.

634 Chaplan SR, Bach FW, Pogrel JW, Chung JM, Yaksh TL (1994) Quantitative assessment of tactile  
635 allodynia in the rat paw. *J Neurosci Methods* 53:55-63.

636 Coggeshall RE, Tate S, Carlton SM (2004) Differential expression of tetrodotoxin-resistant sodium  
637 channels Nav1.8 and Nav1.9 in normal and inflamed rats. *NeurosciLett* 355:45-48.

638 De Col R, Messlinger K, Carr RW (2008) Conduction velocity is regulated by sodium channel  
639 inactivation in unmyelinated axons innervating the rat cranial meninges. *JPhysiol* 586:1089-  
640 1103.

641 De Col R, Messlinger K, Carr RW (2012) Repetitive activity slows axonal conduction velocity and  
642 concomitantly increases mechanical activation threshold in single axons of the rat cranial  
643 dura. *JPhysiol* 590:725-736.

644 Dickie AC, Torsney C (2014) The chemerin receptor 23 agonist, chemerin, attenuates monosynaptic  
645 C-fibre input to lamina I neurokinin 1 receptor expressing rat spinal cord neurons in  
646 inflammatory pain. *MolPain* 10:24.

647 Djouhri L, Lawson SN (1999) Changes in somatic action potential shape in guinea-pig nociceptive  
648 primary afferent neurones during inflammation in vivo. *J Physiol* 520 Pt 2:565-576.

649 Djouhri L, Lawson SN (2001) Increased conduction velocity of nociceptive primary afferent neurons  
650 during unilateral hindlimb inflammation in the anaesthetised guinea-pig. *Neuroscience*  
651 102:669-679.

652 Djouhri L, Koutsikou S, Fang X, McMullan S, Lawson SN (2006) Spontaneous pain, both neuropathic  
653 and inflammatory, is related to frequency of spontaneous firing in intact C-fiber nociceptors.  
654 *JNeurosci* 26:1281-1292.

655 Djouhri L, Al Otaibi M, Kahlat K, Smith T, Sathish J, Weng X (2015) Persistent hindlimb inflammation  
656 induces changes in activation properties of hyperpolarization-activated current (I<sub>h</sub>) in rat C-  
657 fiber nociceptors in vivo. *Neuroscience* 301:121-133.

658 Emery EC, Young GT, McNaughton PA (2012) HCN2 ion channels: an emerging role as the  
659 pacemakers of pain. *Trends PharmacolSci* 33:456-463.

660 Emery EC, Young GT, Berrocoso EM, Chen L, McNaughton PA (2011) HCN2 ion channels play a  
661 central role in inflammatory and neuropathic pain. *Science* 333:1462-1466.

662 Flake NM, Bonebreak DB, Gold MS (2005) Estrogen and inflammation increase the excitability of rat  
663 temporomandibular joint afferent neurons. *Journal of neurophysiology* 93:1585-1597.

664 Gee MD, Lynn B, Cotsell B (1996) Activity-dependent slowing of conduction velocity provides a  
665 method for identifying different functional classes of C-fibre in the rat saphenous nerve.  
666 *Neuroscience* 73:667-675.

667 Gold MS, Gebhart GF (2010) Nociceptor sensitization in pain pathogenesis. *Nat Med* 16:1248-1257.

668 Gould HJ, III, England JD, Soignier RD, Nolan P, Minor LD, Liu ZP, Levinson SR, Paul D (2004)  
669 Ibuprofen blocks changes in Na v 1.7 and 1.8 sodium channels associated with complete  
670 Freund's adjuvant-induced inflammation in rat. *JPain* 5:270-280.

671 Hoffmann T, Kistner K, Nassar M, Reeh PW, Weidner C (2016) Use dependence of peripheral  
672 nociceptive conduction in the absence of tetrodotoxin-resistant sodium channel subtypes. *J*  
673 *Physiol* 594:5529-5541.

674 Ikeda H, Heinke B, Ruscheweyh R, Sandkuhler J (2003) Synaptic plasticity in spinal lamina I projection  
675 neurons that mediate hyperalgesia. *Science* 299:1237-1240.

676 Ikeda H, Stark J, Fischer H, Wagner M, Drdla R, Jager T, Sandkuhler J (2006) Synaptic amplifier of  
677 inflammatory pain in the spinal dorsal horn. *Science* 312:1659-1662.

678 Kim YS, Anderson M, Park K, Zheng Q, Agarwal A, Gong C, Saijilafu, Young L, He S, LaVinka PC, Zhou F,  
679 Bergles D, Hanani M, Guan Y, Spray DC, Dong X (2016) Coupled Activation of Primary Sensory  
680 Neurons Contributes to Chronic Pain. *Neuron* 91:1085-1096.

681 Kist AM et al. (2016) SCN10A Mutation in a Patient with Erythromelalgia Enhances C-Fiber Activity  
682 Dependent Slowing. *PloS one* 11:e0161789.

683 Kleggetveit IP, Namer B, Schmidt R, Helas T, Ruckel M, Orstavik K, Schmelz M, Jorum E (2012) High  
684 spontaneous activity of C-nociceptors in painful polyneuropathy. *Pain* 153:2040-2047.

685 Labrakakis C, MacDermott AB (2003) Neurokinin receptor 1-expressing spinal cord neurons in lamina  
686 I and III/IV of postnatal rats receive inputs from capsaicin sensitive fibers. *Neuroscience*  
687 letters 352:121-124.

688 Lee JH, Park CK, Chen G, Han Q, Xie RG, Liu T, Ji RR, Lee SY (2014) A monoclonal antibody that targets  
689 a Nav1.7 channel voltage sensor for pain and itch relief. *Cell* 157:1393-1404.

690 Leem JW, Willis WD, Chung JM (1993) Cutaneous sensory receptors in the rat foot. *Journal of*  
691 *neurophysiology* 69:1684-1699.

692 Li J, Baccei ML (2016) Neonatal Tissue Damage Promotes Spike Timing-Dependent Synaptic Long-  
693 Term Potentiation in Adult Spinal Projection Neurons. *The Journal of neuroscience : the*  
694 *official journal of the Society for Neuroscience* 36:5405-5416.

695 Liang L, Fan L, Tao B, Yaster M, Tao YX (2013) Protein kinase B/Akt is required for complete Freund's  
696 adjuvant-induced upregulation of Nav1.7 and Nav1.8 in primary sensory neurons. *The*  
697 *journal of pain : official journal of the American Pain Society* 14:638-647.

698 Lynn B, Carpenter SE (1982) Primary afferent units from the hairy skin of the rat hind limb. *Brain Res*  
699 238:29-43.

700 Mantyh PW, Rogers SD, Honore P, Allen BJ, Ghilardi JR, Li J, Daughters RS, Lappi DA, Wiley RG,  
701 Simone DA (1997) Inhibition of hyperalgesia by ablation of lamina I spinal neurons  
702 expressing the substance P receptor. *Science* 278:275-279.

703 Marshall GE, Shehab SA, Spike RC, Todd AJ (1996) Neurokinin-1 receptors on lumbar spinothalamic  
704 neurons in the rat. *Neuroscience* 72:255-263.

705 Matson DJ, Hamamoto DT, Bregman H, Cooke M, DiMauro EF, Huang L, Johnson D, Li X, McDermott  
706 J, Morgan C, Wilenkin B, Malmberg AB, McDonough SI, Simone DA (2015) Inhibition of  
707 Inactive States of Tetrodotoxin-Sensitive Sodium Channels Reduces Spontaneous Firing of C-  
708 Fiber Nociceptors and Produces Analgesia in Formalin and Complete Freund's Adjuvant  
709 Models of Pain. *PloS one* 10:e0138140.

710 Mazo I, Rivera-Arconada I, Roza C (2013) Axotomy-induced changes in activity-dependent slowing in  
 711 peripheral nerve fibres: role of hyperpolarization-activated/HCN channel current. *EurJPain*  
 712 17:1281-1290.

713 McGowan E, Hoyt SB, Li X, Lyons KA, Abbadie C (2009) A peripherally acting Na(v)1.7 sodium channel  
 714 blocker reverses hyperalgesia and allodynia on rat models of inflammatory and neuropathic  
 715 pain. *AnesthAnalg* 109:951-958.

716 Mogil JS (2012) Sex differences in pain and pain inhibition: multiple explanations of a controversial  
 717 phenomenon. *NatRevNeurosci* 13:859-866.

718 Mogil JS, Bailey AL (2010) Sex and gender differences in pain and analgesia. *ProgBrain Res* 186:141-  
 719 157.

720 Nakatsuka T, Ataka T, Kumamoto E, Tamaki T, Yoshimura M (2000) Alteration in synaptic inputs  
 721 through C-afferent fibers to substantia gelatinosa neurons of the rat spinal dorsal horn  
 722 during postnatal development. *Neuroscience* 99:549-556.

723 Namer B, Orstavik K, Schmidt R, Kleggetveit IP, Weidner C, Mork C, Kvernebo MS, Kvernebo K, Salter  
 724 H, Carr TH, Segerdahl M, Quiding H, Waxman SG, Handwerker HO, Torebjork HE, Jorum E,  
 725 Schmelz M (2015) Specific changes in conduction velocity recovery cycles of single  
 726 nociceptors in a patient with erythromelalgia with the I848T gain-of-function mutation of  
 727 Nav1.7. *Pain* 156:1637-1646.

728 Nassar MA, Levato A, Stirling LC, Wood JN (2005) Neuropathic pain develops normally in mice lacking  
 729 both Na(v)1.7 and Na(v)1.8. *MolPain* 1:24.

730 Nassar MA, Stirling LC, Forlani G, Baker MD, Matthews EA, Dickenson AH, Wood JN (2004)  
 731 Nociceptor-specific gene deletion reveals a major role for Nav1.7 (PN1) in acute and  
 732 inflammatory pain. *ProcNatlAcadSciUSA* 101:12706-12711.

733 Nichols ML, Allen BJ, Rogers SD, Ghilardi JR, Honore P, Luger NM, Finke MP, Li J, Lappi DA, Simone  
 734 DA, Mantyh PW (1999) Transmission of chronic nociception by spinal neurons expressing the  
 735 substance P receptor. *Science* 286:1558-1561.

736 Obreja O, Hirth M, Turnquist B, Rukwied R, Ringkamp M, Schmelz M (2012) The differential effects of  
 737 two sodium channel modulators on the conductive properties of C-fibers in pig skin in vivo.  
 738 *Anesth Analg* 115:560-571.

739 Obreja O, Ringkamp M, Namer B, Forsch E, Klusch A, Rukwied R, Petersen M, Schmelz M (2010)  
 740 Patterns of activity-dependent conduction velocity changes differentiate classes of  
 741 unmyelinated mechano-insensitive afferents including cold nociceptors, in pig and in human.  
 742 *Pain* 148:59-69.

743 Obreja O, Ringkamp M, Turnquist B, Hirth M, Forsch E, Rukwied R, Petersen M, Schmelz M (2011)  
 744 Nerve growth factor selectively decreases activity-dependent conduction slowing in  
 745 mechano-insensitive C-nociceptors. *Pain* 152:2138-2146.

746 Orstavik K, Weidner C, Schmidt R, Schmelz M, Hilliges M, Jorum E, Handwerker H, Torebjork E (2003)  
 747 Pathological C-fibres in patients with a chronic painful condition. *Brain* 126:567-578.

748 Orstavik K, Namer B, Schmidt R, Schmelz M, Hilliges M, Weidner C, Carr RW, Handwerker H, Jorum E,  
 749 Torebjork HE (2006) Abnormal function of C-fibers in patients with diabetic neuropathy. *The*  
 750 *Journal of neuroscience : the official journal of the Society for Neuroscience* 26:11287-  
 751 11294.

752 Otsuka M, Konishi S (1974) Electrophysiology of mammalian spinal cord in vitro. *Nature* 252:733-  
 753 734.

754 Papp I, Hollo K, Antal M (2010) Plasticity of hyperpolarization-activated and cyclic nucleotid-gated  
 755 cation channel subunit 2 expression in the spinal dorsal horn in inflammatory pain.  
 756 *EurJNeurosci* 32:1193-1201.

757 Payne CE et al. (2015) A novel selective and orally bioavailable Nav 1.8 channel blocker, PF-  
 758 01247324, attenuates nociception and sensory neuron excitability. *Br J Pharmacol* 172:2654-  
 759 2670.

760 Peirs C, Williams SP, Zhao X, Walsh CE, Gedeon JY, Cagle NE, Goldring AC, Hioki H, Liu Z, Marell PS,  
 761 Seal RP (2015) Dorsal Horn Circuits for Persistent Mechanical Pain. *Neuron* 87:797-812.



762 Petersson ME, Obreja O, Lampert A, Carr RW, Schmelz M, Fransen E (2014) Differential axonal  
 763 conduction patterns of mechano-sensitive and mechano-insensitive nociceptors - a  
 764 combined experimental and modelling study. PLoSOne 9:e103556.  
 765 Rahman W, Dickenson AH (2013) Voltage gated sodium and calcium channel blockers for the  
 766 treatment of chronic inflammatory pain. NeurosciLett 557 Pt A:19-26.  
 767 Rang HP, Ritchie JM (1968) On the electrogenic sodium pump in mammalian non-myelinated nerve  
 768 fibres and its activation by various external cations. J Physiol 196:183-221.  
 769 Scheff NN, Gold MS (2011) Sex differences in the inflammatory mediator-induced sensitization of  
 770 dural afferents. Journal of neurophysiology 106:1662-1668.  
 771 Schmelz M, Schmidt R, Bickel A, Handwerker HO, Torebjork HE (1997) Specific C-receptors for itch in  
 772 human skin. JNeurosci 17:8003-8008.  
 773 Schmelz M, Forster C, Schmidt R, Ringkamp M, Handwerker HO, Torebjork HE (1995) Delayed  
 774 responses to electrical stimuli reflect C-fiber responsiveness in human microneurography.  
 775 Exp Brain Res 104:331-336.  
 776 Schmidt R, Schmelz M, Forster C, Ringkamp M, Torebjork E, Handwerker H (1995) Novel classes of  
 777 responsive and unresponsive C nociceptors in human skin. JNeurosci 15:333-341.  
 778 Serra J, Campero M, Ochoa J, Bostock H (1999) Activity-dependent slowing of conduction  
 779 differentiates functional subtypes of C fibres innervating human skin. J Physiol 515 ( Pt  
 780 3):799-811.  
 781 Serra J, Collado A, Sola R, Antonelli F, Torres X, Salgueiro M, Quiles C, Bostock H (2014)  
 782 Hyperexcitable C nociceptors in fibromyalgia. Ann Neurol 75:196-208.  
 783 Shim B, Ringkamp M, Lambrinos GL, Hartke TV, Griffin JW, Meyer RA (2007) Activity-dependent  
 784 slowing of conduction velocity in uninjured L4 C fibers increases after an L5 spinal nerve  
 785 injury in the rat. Pain 128:40-51.  
 786 Spike RC, Puskar Z, Andrew D, Todd AJ (2003) A quantitative and morphological study of projection  
 787 neurons in lamina I of the rat lumbar spinal cord. EurJNeurosci 18:2433-2448.

788 Sun W, Miao B, Wang XC, Duan JH, Wang WT, Kuang F, Xie RG, Xing JL, Xu H, Song XJ, Luo C, Hu SJ  
 789 (2012) Reduced conduction failure of the main axon of polymodal nociceptive C-fibres  
 790 contributes to painful diabetic neuropathy in rats. *Brain* 135:359-375.

791 Suzuki R, Morcuende S, Webber M, Hunt SP, Dickenson AH (2002) Superficial NK1-expressing  
 792 neurons control spinal excitability through activation of descending pathways. *Nature*  
 793 *neuroscience* 5:1319-1326.

794 Takigawa T, Alzheimer C, Quasthoff S, Grafe P (1998) A special blocker reveals the presence and  
 795 function of the hyperpolarization-activated cation current  $I_H$  in peripheral mammalian nerve  
 796 fibres. *Neuroscience* 82:631-634.

797 Thalhammer JG, Raymond SA, Popitz-Bergez FA, Strichartz GR (1994) Modality-dependent  
 798 modulation of conduction by impulse activity in functionally characterized single cutaneous  
 799 afferents in the rat. *Somatosens Mot Res* 11:243-257.

800 Thompson SW, Dray A, Urban L (1994) Injury-induced plasticity of spinal reflex activity: NK1  
 801 neurokinin receptor activation and enhanced A- and C-fiber mediated responses in the rat  
 802 spinal cord in vitro. *The Journal of neuroscience : the official journal of the Society for*  
 803 *Neuroscience* 14:3672-3687.

804 Thompson SW, Gerber G, Sivilotti LG, Woolf CJ (1992) Long duration ventral root potentials in the  
 805 neonatal rat spinal cord in vitro; the effects of ionotropic and metabotropic excitatory amino  
 806 acid receptor antagonists. *Brain Res* 595:87-97.

807 Tigerholm J, Petersson ME, Obreja O, Lampert A, Carr R, Schmelz M, Fransen E (2014) Modeling  
 808 activity-dependent changes of axonal spike conduction in primary afferent C-nociceptors.  
 809 *JNeurophysiol* 111:1721-1735.

810 Todd AJ, McGill MM, Shehab SA (2000) Neurokinin 1 receptor expression by neurons in laminae I, III  
 811 and IV of the rat spinal dorsal horn that project to the brainstem. *EurJNeurosci* 12:689-700.

812 Tong CK, MacDermott AB (2006) Both  $Ca^{2+}$ -permeable and -impermeable AMPA receptors  
 813 contribute to primary synaptic drive onto rat dorsal horn neurons. *J Physiol* 575:133-144.

814 Torebjork HE, LaMotte RH, Robinson CJ (1984) Peripheral neural correlates of magnitude of  
815 cutaneous pain and hyperalgesia: simultaneous recordings in humans of sensory judgments  
816 of pain and evoked responses in nociceptors with C-fibers. *Journal of neurophysiology*  
817 51:325-339.

818 Torsney C (2011) Inflammatory pain unmasks heterosynaptic facilitation in lamina I neurokinin 1  
819 receptor-expressing neurons in rat spinal cord. *JNeurosci* 31:5158-5168.

820 Torsney C, MacDermott AB (2006) Disinhibition opens the gate to pathological pain signaling in  
821 superficial neurokinin 1 receptor-expressing neurons in rat spinal cord. *JNeurosci* 26:1833-  
822 1843.

823 Wang F, Cai B, Li KC, Hu XY, Lu YJ, Wang Q, Bao L, Zhang X (2015) FXD2, a gamma subunit of Na(+),  
824 K(+)-ATPase, maintains persistent mechanical allodynia induced by inflammation. *Cell Res*  
825 25:318-334.

826 Wang X, Wang S, Wang W, Duan J, Zhang M, Lv X, Niu C, Tan C, Wu Y, Yang J, Hu S, Xing J (2016a) A  
827 novel intrinsic analgesic mechanism: the enhancement of the conduction failure along  
828 polymodal nociceptive C-fibers. *Pain* 157:2235-2247.

829 Wang XC, Wang S, Zhang M, Gao F, Yin C, Li H, Zhang Y, Hu SJ, Duan JH (2016b) Alpha-Dendrotoxin-  
830 sensitive Kv1 channels contribute to conduction failure of polymodal nociceptive C-fibers  
831 from rat coccygeal nerve. *Journal of neurophysiology* 115:947-957.

832 Waxman SG, Zamponi GW (2014) Regulating excitability of peripheral afferents: emerging ion  
833 channel targets. *NatNeurosci* 17:153-163.

834 Weidner C, Schmelz M, Schmidt R, Hansson B, Handwerker HO, Torebjork HE (1999) Functional  
835 attributes discriminating mechano-insensitive and mechano-responsive C nociceptors in  
836 human skin. *The Journal of neuroscience : the official journal of the Society for Neuroscience*  
837 19:10184-10190.

838 Weidner C, Schmelz M, Schmidt R, Hammarberg B, Orstavik K, Hilliges M, Torebjork HE, Handwerker  
839 HO (2002) Neural signal processing: the underestimated contribution of peripheral human C-

840 fibers. The Journal of neuroscience : the official journal of the Society for Neuroscience  
841 22:6704-6712.

842 Weng X, Smith T, Sathish J, Djouhri L (2012) Chronic inflammatory pain is associated with increased  
843 excitability and hyperpolarization-activated current (I<sub>h</sub>) in C- but not A-delta-nociceptors.  
844 Pain 153:900-914.

845 Xiao WH, Bennett GJ (2007) Persistent low-frequency spontaneous discharge in A-fiber and C-fiber  
846 primary afferent neurons during an inflammatory pain condition. Anesthesiology 107:813-  
847 821.

848 Yang S, Xiao Y, Kang D, Liu J, Li Y, Undheim EA, Klint JK, Rong M, Lai R, King GF (2013) Discovery of a  
849 selective NaV1.7 inhibitor from centipede venom with analgesic efficacy exceeding  
850 morphine in rodent pain models. Proc Natl Acad Sci U S A 110:17534-17539.

851 Yarnitsky D, Ochoa JL (1990) Studies of heat pain sensation in man: perception thresholds, rate of  
852 stimulus rise and reaction time. Pain 40:85-91.

853 Young GT, Emery EC, Mooney ER, Tsantoulas C, McNaughton PA (2014) Inflammatory and  
854 neuropathic pain are rapidly suppressed by peripheral block of hyperpolarisation-activated  
855 cyclic nucleotide-gated ion channels. Pain 155:1708-1719.

856 Zhang XF, Shieh CC, Chapman ML, Matulenko MA, Hakeem AH, Atkinson RN, Kort ME, Marron BE,  
857 Joshi S, Honore P, Faltynek CR, Krafte DS, Jarvis MF (2010) A-887826 is a structurally novel,  
858 potent and voltage-dependent Na(v)1.8 sodium channel blocker that attenuates neuropathic  
859 tactile allodynia in rats. Neuropharmacology 59:201-207.

860 Zhang XL, Mok LP, Lee KY, Charbonnet M, Gold MS (2012) Inflammation-induced changes in BK(Ca)  
861 currents in cutaneous dorsal root ganglion neurons from the adult rat. Mol Pain 8:37.

862 Zhang YH, Nicol GD (2004) NGF-mediated sensitization of the excitability of rat sensory neurons is  
863 prevented by a blocking antibody to the p75 neurotrophin receptor. Neuroscience letters  
864 366:187-192.

865 Zhu ZR, Tang XW, Wang WT, Ren W, Xing JL, Zhang JR, Duan JH, Wang YY, Jiao X, Hu SJ (2009)  
866 Conduction failures in rabbit saphenous nerve unmyelinated fibers. *Neurosignals* 17:181-  
867 195.

868

## 869 Figure Legends

870 *Figure 1*

871 CFA inflammation reduces C fibre activity-dependent slowing (ADS) in isolated  
872 dorsal roots. **Ai** Two suction electrodes were used to stimulate and record compound  
873 action potentials from L4/L5 dorsal roots. Representative compound action potentials  
874 recorded from dorsal roots isolated from control (**Aii**) and CFA (**Aiii**) treated rats,  
875 illustrating the fast ( $A\beta$ ), medium ( $A\delta$ ) and slow (C) conducting components. The x16  
876 traces recorded in response to 2Hz dorsal root stimulation are shown. Arrowheads  
877 indicate the stimulus artefact. Inserts magnify the C fibre component with broken  
878 lines indicating the negative peaks of the first and last responses. Repetitive  
879 stimulation of dorsal roots at 2Hz (**B**) results in a negligible change in the latency of  
880  $A\beta$  (**i**) or  $A\delta$  fibre (**ii**) responses whereas C fibres display a progressive increase in  
881 response latency (**iii**). **C** AUC analysis of latency change reveals that CFA  
882 inflammation reduces the frequency dependent progressive latency change  
883 observed in C fibres (2-way ANOVA, CFA  $p<0.0001$ ; frequency  $p<0.0001$ ). **D**  
884 Eliminating the contribution of action potential initiation time to latency change, by  
885 subtracting the latency values obtained at the short (position 2) from the long  
886 (position 1) root stimulation length (**i**) confirms that CFA attenuates C fibre ADS (**ii**,  
887 **iii**) (2-way ANOVA, CFA  $p<0.0001$ ). Data in **B**, **C**:  $A\beta$  control  $n=10$ , CFA  $n=12$ ;  $A\delta$ /C  
888 control  $n=11$ , CFA  $n=13$ . Data in **D**; control  $n=10$ , CFA  $n=12$ .

889

890 *Figure 2*

891 CFA inflammation does not alter limited ADS in monosynaptic A $\delta$  fibre input to  
892 lamina I NK1R+ neurons. **Ai** eEPSCs were recorded from pre-labelled lamina I  
893 NK1R+ neurons, in spinal cord slices in response to stimulation of attached dorsal  
894 roots. Representative monosynaptic A $\delta$  fibre eEPSCs recorded in tissue isolated  
895 from control (**Aii**) and CFA (**Aiii**) treated rats. Each trace comprises 16 traces  
896 recorded in response to dorsal root stimulation at 2Hz. Broken lines and arrows  
897 denote the latency of the first and last trace, measured as the time between the  
898 stimulus artefact (denoted by the arrowhead) and the onset of the monosynaptic  
899 response. **B** Stimulation of monosynaptic A $\delta$  fibre input to lamina I NK1R+ neurons  
900 at 1 (i) or 2Hz (ii) results in a small degree of ADS. **C** AUC analysis of latency  
901 change reveals that CFA inflammation does not affect the small frequency  
902 dependent progressive latency change (2-way ANOVA, CFA  $p=0.570$ ; frequency  
903  $p=0.016$ ). 1Hz: control  $n=7$ , CFA  $n=12$ ; 2Hz: control  $n=13$ , CFA  $n=22$ . Note that  
904 scaling in B and C is identical to that in Figure 3.

905

906 *Figure 3*

907 CFA inflammation reduces ADS in monosynaptic C fibre input to lamina I NK1R+  
908 neurons. **A** Representative monosynaptic C fibre eEPSCs recorded in tissue isolated  
909 from control (i) and CFA (ii) treated rats. Each trace comprises 16 traces recorded in  
910 response to repetitive dorsal root stimulation at 1Hz. Broken lines and arrows denote  
911 the latency of the first and last trace, measured as the time between the stimulus  
912 artefact (denoted by the arrowhead) and the onset of the monosynaptic response. **B**  
913 Stimulation of monosynaptic C fibre input to lamina I NK1R+ neurons at 1 (i) or 2Hz

(ii) results in a progressive increase in response latency. AUC analysis of latency change (**C**) reveals that CFA attenuates ADS but it is not affected by stimulation frequency (2-way ANOVA, CFA  $p=0.013$ ; frequency  $p=0.521$ ). 1Hz: control  $n=42$ , CFA  $n=64$ ; 2Hz: control  $n=16$ , CFA  $n=36$ .

*Figure 4*

CFA inflammation reduces the average and range of ADS within the C fibre population. **A** C fibre compound action potentials recorded in response to trains of 40 stimuli delivered at 1 (i) or 2Hz (ii) results in a progressive slowing of the response latency. AUC analysis of latency change (**iii**) reveals that CFA inflammation reduces the frequency dependent progressive latency change (2-way ANOVA, CFA  $p=0.005$ ; frequency  $p<0.0001$ ). **B** Representative C fibre compound action potentials recorded, in response to 2Hz stimulation, from dorsal roots isolated from control (i) and CFA (ii) treated rats, with broken lines and arrows indicating the width (positive-positive peak) of the first and last response. **C** Trains of 40 stimuli delivered at 1 (i) or 2Hz (ii) result in a progressive increase in C fibre compound action potential width. AUC analysis of width change (**iii**) reveals that CFA inflammation reduces the frequency dependent progressive width increase (2-way ANOVA, CFA  $p<0.0001$ ; frequency  $p<0.0001$ ). All: control  $n=18$ , CFA  $n=16$ .

*Figure 5*

CFA inflammation reduces ADS in monosynaptic C fibre input to lamina I NK1R+ neurons during high frequency stimulus trains but does not alter synaptic response failures. **A**. Repetitive stimulation of monosynaptic C fibre input to lamina I NK1R+ neurons, using trains of 40 stimuli, delivered at 2 (i), 5 (ii) or 10Hz (iii) results (iv) in

a frequency dependent progressive increase in the eEPSC latency, that is significantly reduced by CFA inflammation (2-way ANOVA, CFA  $p<0.0001$ ; frequency  $p=0.039$ ). **B.** There is a progressive increase in the number of synaptic response failures per stimulus number during stimulation of monosynaptic C fibre input to lamina I NK1R+ neurons at 2 (i), 5 (ii) or 10Hz (iii). The total number of synaptic response failures (iv) is frequency dependent but not altered by CFA inflammation (2-way RM ANOVA, CFA  $p=0.971$ ; frequency  $p<0.0001$ ). All groups  $n=9$ .

*Figure 6*

CFA inflammation reduces ADS, within the C fibre population and in the monosynaptic C fibre input to lamina I NK1R+ neurons, in females only. C fibre compound action potentials recorded in response to 2Hz trains (x40 stimuli) in females (A) and males (B) results in a progressive slowing of the response latency (i) and width (ii). C AUC analysis of latency change (i) and width change (ii) reveals a significant interaction between sex and inflammation (2-way ANOVA latency change  $p=0.037$ , width change  $p=0.003$ ) with CFA inflammation reducing ADS in females only (latency change females  $p=0.031$ , males  $p=0.998$ ; width change females  $p<0.0001$ , males  $p=0.525$ , Tukey multiple comparisons test). Monosynaptic eEPSCs in lamina I NK1R+ neurons recorded in response to 2Hz trains (x40 stimuli) results in a progressive slowing of the response latency in females (D) and males (E). F AUC analysis of latency change reveals a significant interaction between sex and inflammation (2-way ANOVA  $p=0.034$ ) with CFA inflammation reducing ADS in females only (female  $p=0.015$ ; male  $p=0.996$ , Tukey multiple comparisons test). CAP: female control  $n=9$ , female CFA  $n=7$ , male control  $n=9$ , male CFA  $n=9$ . eEPSC: female control  $n=10$ , female CFA  $n=17$ , male control  $n=6$ , male CFA  $n=16$ .



*Figure 7*

Cumulative dorsal root evoked ventral root potentials are regulated in a frequency and length dependent manner. **A** Ventral root potentials were recorded from a hemisected spinal cord preparation in response to long length and short length dorsal root stimulation (x20) to increase and decrease the spinal impact of ADS, respectively, using suction electrodes. **B** Representative cumulative dorsal root evoked (5Hz) ventral root potential **(i)** that is abolished in the presence of NBQX (10 $\mu$ M) and APV (50 $\mu$ M) **(ii)**. Stimulus artefact visible as vertical lines in all traces. **C** Representative cumulative dorsal root evoked ventral root potentials induced by 2, 5 and 10Hz stimulation over long **(i)** and short **(ii)** dorsal root lengths. Arrows denote the 500ms timepoint, following the last stimulus artefact, when amplitude is measured. **D(i)** Repeated measures ANOVA of normalised cumulative ventral root potentials reveals both frequency ( $p=0.039$ ) and length ( $p=0.003$ ) dependent effects that significantly interact ( $p=0.022$ ) with Sidak's multiple comparisons test revealing a significant impact of length at 2Hz only ( $p<0.01$ ),  $n=10$ . **(ii)** Difference in % cumulative depolarisation between long and short stimulation lengths is significantly greater in females (One-tailed Mann-Whitney test,  $p=0.028$ ),  $n=5$  both groups.

*Figure 8*

Behavioural analysis of the nociceptive flexion withdrawal reflex demonstrates elevated thermal pain thresholds and enhanced inflammatory thermal hyperalgesia in females. **A** Ipsilateral and contralateral hindpaw withdrawal latencies to noxious radiant heat stimuli (Hargreaves) were measured in female and male rats prior to and 2-5d following CFA intraplantar injection. **B** Ipsilateral withdrawal latency values expressed as a percentage of contralateral values, reveal a sex difference in thermal

989 hyperalgesia (repeated measures-ANOVA,  $p=0.032$ ). **C** AUC analysis (2-5d post  
990 CFA) shows a significant sex difference in contralateral hindpaw withdrawal latency  
991 with higher values in females (2-way ANOVA followed by Sidak's multiple  
992 comparisons test,  $p=0.022$ ). **D** Ipsilateral hindpaw thickness is expressed as a  
993 percentage of contralateral values and demonstrates that CFA-induced paw swelling  
994 is not sex dependent (repeated measures ANOVA,  $p=0.513$ ). **E** Ipsilateral and  
995 contralateral hindpaw mechanical thresholds (von Frey) were also measured in the  
996 same female and male rats prior to and 2-5d following CFA intraplantar injection. **F**  
997 Ipsilateral mechanical thresholds are expressed as a percentage of contralateral  
998 values and demonstrates that mechanical inflammatory hypersensitivity is not sex  
999 dependent (repeated measures ANOVA,  $p=0.548$ ). All graphs female  $n=9$ , male  $n=7$ .